



TITLE:

# N-Fractional Calculus of Some Multi-Powers Functions(Study on Geometric Univalent Function Theory)

AUTHOR(S):

Nishimoto, Katsuyuki

---

CITATION:

Nishimoto, Katsuyuki. N-Fractional Calculus of Some Multi-Powers Functions(Study on Geometric Univalent Function Theory). 数理解析研究所講究録 2008, 1579: 79-87

ISSUE DATE:

2008-02

URL:

<http://hdl.handle.net/2433/81390>

RIGHT:

## N- Fractional Calculus of Some Multi- Powers Functions

Katsuyuki Nishimoto

*Institute for Applied Mathematics, Descartes Press Co.*

*2- 13- 10 Kaguike, Koriyama, 963 - 8833, JAPAN*

*Fax : + 81 - 24 - 922 - 7596*

### Abstract

In a previous article of the author, N- fractional calculus

$$\left( ((z-b)^\beta - c)^\alpha \right)_\nu, \quad ((z-b)^\beta - c \neq 0)$$

are discussed. In this article that of more extended forms

$$\left( (((z-b)^\beta - c)^\alpha - d)^\delta \right)_\nu, \quad (((z-b)^\beta - c)^\alpha - d \neq 0)$$

are discussed. Moreover their special cases

$$\left( (((z-b)^\beta - c)^\alpha - d)^\delta \right)_n, \quad (n \in \mathbb{Z}_0^+, ((z-b)^\beta - c)^\alpha - d \neq 0)$$

are presented .

### § 0. Introduction ( Definition of Fractional Calculus )

( I ) Definition. ( by K. Nishimoto ) ( [ 1 ] Vol. 1 )

Let  $D = \{D_-, D_+\}$ ,  $C = \{C_-, C_+\}$ ,

$C_-$  be a curve along the cut joining two points  $z$  and  $-\infty + i\text{Im}(z)$ ,

$C_+$  be a curve along the cut joining two points  $z$  and  $\infty + i\text{Im}(z)$ ,

$D_-$  be a domain surrounded by  $C_-$ ,  $D_+$  be a domain surrounded by  $C_+$  .

( Here  $D$  contains the points over the curve  $C$  ).

Moreover, let  $f = f(z)$  be a regular function in  $D(z \in D)$ ,

$$f_\nu = (f)_\nu = {}_c(f)_\nu = \frac{\Gamma(\nu+1)}{2\pi i} \int_c \frac{f(\xi)}{(\xi-z)^{\nu+1}} d\xi \quad (\nu \notin \mathbb{Z}), \quad (1)$$

$$(f)_{-m} = \lim_{\nu \rightarrow -m} (f)_\nu \quad (m \in \mathbb{Z}^+), \quad (2)$$

where  $-\pi \leq \arg(\xi-z) \leq \pi$  for  $C_-$ ,  $0 \leq \arg(\xi-z) \leq 2\pi$  for  $C_+$ ,

$\xi \neq z$ ,  $z \in C$ ,  $\nu \in \mathbb{R}$ ,  $\Gamma$ ; Gamma function,

then  $(f)_\nu$  is the fractional differintegration of arbitrary order  $\nu$  ( derivatives of order  $\nu$  for  $\nu > 0$ , and integrals of order  $-\nu$  for  $\nu < 0$  ), with respect to  $z$ , of the function  $f$ , if  $|(f)_\nu| < \infty$ .

(II) On the fractional calculus operator  $N^\nu$  [3]

**Theorem A.** Let fractional calculus operator (Nishimoto's Operator)  $N^\nu$  be

$$N^\nu = \left( \frac{\Gamma(\nu+1)}{2\pi i} \int_C \frac{d\xi}{(\xi-z)^{\nu+1}} \right) \quad (\nu \notin \mathbb{Z}), \quad [\text{Refer to (1)}] \quad (3)$$

with 
$$N^{-m} = \lim_{\nu \rightarrow -m} N^\nu \quad (m \in \mathbb{Z}^+), \quad (4)$$

and define the binary operation  $\circ$  as

$$N^\beta \circ N^\alpha f = N^\beta N^\alpha f = N^\beta(N^\alpha f) \quad (\alpha, \beta \in \mathbb{R}), \quad (5)$$

then the set

$$\{N^\nu\} = \{N^\nu | \nu \in \mathbb{R}\} \quad (6)$$

is an Abelian product group (having continuous index  $\nu$ ) which has the inverse transform operator  $(N^\nu)^{-1} = N^{-\nu}$  to the fractional calculus operator  $N^\nu$ , for the function  $f$  such that  $f \in F = \{f; 0 \neq |f_\nu| < \infty, \nu \in \mathbb{R}\}$ , where  $f = f(z)$  and  $z \in C$ . (vis.  $-\infty < \nu < \infty$ ).

(For our convenience, we call  $N^\beta \circ N^\alpha$  as product of  $N^\beta$  and  $N^\alpha$ .)

**Theorem B.** "F.O.G.  $\{N^\nu\}$ " is an "Action product group which has continuous index  $\nu$ " for the set of  $F$ . (F.O.G.; Fractional calculus operator group) [3]

(III) Lemma. We have [1]

$$(i) \quad ((z-c)^b)_\alpha = e^{-i\pi\alpha} \frac{\Gamma(\alpha-b)}{\Gamma(-b)} (z-c)^{b-\alpha} \quad \left( \left| \frac{\Gamma(\alpha-b)}{\Gamma(-b)} \right| < \infty \right), \quad (7)$$

$$(ii) \quad (\log(z-c))_\alpha = -e^{-i\pi\alpha} \Gamma(\alpha) (z-c)^{-\alpha} \quad (|\Gamma(\alpha)| < \infty), \quad (8)$$

$$(iii) \quad ((z-c)^{-\alpha})_{-\alpha} = -e^{i\pi\alpha} \frac{1}{\Gamma(\alpha)} \log(z-c) \quad (|\Gamma(\alpha)| < \infty), \quad (9)$$

where  $z-c \neq 0$  for (i) and  $z-c \neq 0, 1$  for (ii), (iii),

$$(iv) \quad (u \cdot v)_\alpha = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha+1)}{k! \Gamma(\alpha+1-k)} u_{\alpha-k} v_k \quad \left( \begin{array}{l} u = u(z), \\ v = v(z) \end{array} \right). \quad (10)$$

### § 1. Preliminary

The Teorem below is reported by the author already (cf. J. Frac. Calc. Vol. 29, May (2006), pp.35 - 44.) . [12]

**Theorem D.** We have

$$(i) \quad (((z-b)^\beta - c)^\alpha)_\gamma = e^{-i\pi\gamma} (z-b)^{\alpha\beta-\gamma}$$

$$\times \sum_{k=0}^{\infty} \frac{[-\alpha]_k \Gamma(\beta k - \alpha \beta + \gamma)}{k! \Gamma(\beta k - \alpha \beta)} \left( \frac{c}{(z-b)^\beta} \right)^k \quad (1)$$

$$\left( \left| \frac{\Gamma(\beta k - \alpha \beta + \gamma)}{\Gamma(\beta k - \alpha \beta)} \right| < \infty \right)$$

and

$$(ii) \quad \left( ((z-b)^\beta - c)^\alpha \right)_n = (-1)^n (z-b)^{\alpha \beta - n}$$

$$\times \sum_{k=0}^{\infty} \frac{[-\alpha]_k [\beta k - \alpha \beta]_n}{k!} \left( \frac{c}{(z-b)^\beta} \right)^k \quad (n \in \mathbb{Z}_0^+) \quad (2)$$

where

$$\left| \frac{c}{(z-b)^\beta} \right| < 1,$$

and

$$[\lambda]_k = \lambda(\lambda+1) \cdots (\lambda+k-1) = \Gamma(\lambda+k)/\Gamma(\lambda) \text{ with } [\lambda]_0 = 1,$$

(Notation of Pochhammer).

## § 2. N-Fractional Calculus of Functions

$$\left( ((z-b)^\beta - c)^\alpha - d \right)^\delta$$

**Theorem 1.** We have

$$(i) \quad \left( (((z-b)^\beta - c)^\alpha - d)^\delta \right)_\gamma = e^{-i\pi\gamma} (z-b)^{\alpha \beta \delta - \gamma} \\ \times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k \Gamma(\beta k - \alpha \beta(\delta-m) + \gamma)}{m! \cdot k! \Gamma(\beta k - \alpha \beta(\delta-m))} \left( \frac{c}{(z-b)^\beta} \right)^k \left( \frac{d}{(z-b)^{\alpha \beta}} \right)^m \quad (1)$$

$$\left( \left| \frac{\Gamma(\beta k - \alpha \beta(\delta-m) + \gamma)}{\Gamma(\beta k - \alpha \beta(\delta-m))} \right| < \infty \right)$$

and

$$(ii) \quad \left( (((z-b)^\beta - c)^\alpha - d)^\delta \right)_n = (-1)^n (z-b)^{\alpha \beta \delta - n}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k [\beta k - \alpha \beta(\delta-m)]_n}{m! \cdot k!} \left( \frac{c}{(z-b)^\beta} \right)^k \left( \frac{d}{(z-b)^{\alpha \beta}} \right)^m, \quad (2)$$

where

$$((z-b)^\beta - c)^\alpha - d \neq 0, \quad \left| \frac{c}{(z-b)^\beta} \right| < 1, \quad \left| \frac{d}{(z-b)^{\alpha\beta}} \right| < 1, \quad n \in \mathbb{Z}_0^+.$$

**Proof of (i).** We have

$$(((z-b)^\beta - c)^\alpha - d)^\delta = X^\delta \left(1 - \frac{d}{X}\right)^\delta \quad (X = ((z-b)^\beta - c)^\alpha) \quad (3)$$

$$= X^\delta \sum_{m=0}^{\infty} \frac{[-\delta]_m}{m!} \left(\frac{d}{X}\right)^m \quad (4)$$

$$= \sum_{m=0}^{\infty} \frac{[-\delta]_m (d^m)}{m!} X^{\delta-m} \quad (5)$$

$$= \sum_{m=0}^{\infty} \frac{[-\delta]_m (d^m)}{m!} ((z-b)^\beta - c)^{\alpha(\delta-m)}, \quad (6)$$

hence, operating N-fractional calculus operator  $N^\gamma$  to the both sides of (6), we obtain

$$\begin{aligned} & \left( (((z-b)^\beta - c)^\alpha - d)^\delta \right)_\gamma \\ &= \sum_{m=0}^{\infty} \frac{[-\delta]_m (d^m)}{m!} \left( ((z-b)^\beta - c)^{\alpha(\delta-m)} \right)_\gamma \end{aligned} \quad (7)$$

$$\begin{aligned} &= \sum_{m=0}^{\infty} \frac{[-\delta]_m (d^m)}{m!} \left[ e^{-i\pi\gamma} (z-b)^{\alpha\beta(\delta-m)-\gamma} \right. \\ &\quad \times \sum_{k=0}^{\infty} \frac{[-\alpha(\delta-m)]_k \Gamma(\beta k - \alpha\beta(\delta-m) + \gamma)}{k! \Gamma(\beta k - \alpha\beta(\delta-m))} \left( \frac{c}{(z-b)^\beta} \right)^k \left. \right], \end{aligned} \quad (8)$$

applying Theorem D. (i), under the conditions stated before.

We have then (1) from (8) clearly.

**Proof of (ii).** Set  $\gamma = n$  in (1).

**Note 1.** We use the notations  $\sum_{m,k=0}^{\infty} \cdots = \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \cdots$ , for our convenience.

**Note.2.** When  $d = 0$  and  $\delta = 1$ , Theorem 1 is reduced to Theorem D., clearly.

**Corollary 1.** We have

$$\begin{aligned}
 (i) \quad & \left( ((z^\beta - c)^\alpha - d)^\delta \right)_\gamma = e^{-i\pi\gamma} z^{\alpha\beta\delta - \gamma} \\
 & \times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta - m)]_k \Gamma(\beta k - \alpha\beta(\delta - m) + \gamma)}{m! \cdot k! \Gamma(\beta k - \alpha\beta(\delta - m))} \left( \frac{c}{z^\beta} \right)^k \left( \frac{d}{z^{\alpha\beta}} \right)^m \\
 & \left( \left| \frac{\Gamma(\beta k - \alpha\beta(\delta - m) + \gamma)}{\Gamma(\beta k - \alpha\beta(\delta - m))} \right| < \infty \right)
 \end{aligned} \quad (9)$$

and

$$\begin{aligned}
 (ii) \quad & \left( ((z^\beta - c)^\alpha - d)^\delta \right)_n = (-1)^n z^{\alpha\beta\delta - n} \\
 & \times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta - m)]_k [\beta k - \alpha\beta(\delta - m)]_n}{m! \cdot k!} \left( \frac{c}{z^\beta} \right)^k \left( \frac{d}{z^{\alpha\beta}} \right)^m,
 \end{aligned} \quad (10)$$

where

$$(z^\beta - c)^\alpha - d \neq 0, \quad \left| \frac{c}{z^\beta} \right| < 1, \quad \left| \frac{d}{z^{\alpha\beta}} \right| < 1, \quad n \in \mathbb{Z}_0^+.$$

**Proof.** Set  $b = 0$  in Theorem 1.

### § 3. Some Special Cases

[I] When  $\beta = \alpha = 1$ , we obtain

$$\begin{aligned}
 & \left( ((z - b - c - d)^\delta \right)_\gamma = e^{-i\pi\gamma} (z - b)^{\delta - \gamma} \\
 & \times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [m - \delta]_k \Gamma(k + m - \delta + \gamma)}{m! \cdot k! \Gamma(k + m - \delta)} \left( \frac{c}{z - b} \right)^k \left( \frac{d}{z - b} \right)^m \\
 & \left( \left| \frac{c}{z - b} \right|, \left| \frac{d}{z - b} \right| < 1, \left| \frac{\Gamma(k + m - \delta + \gamma)}{\Gamma(k + m - \delta)} \right| < \infty \right)
 \end{aligned} \quad (1)$$

from Theorem 1. (i).

Now we have the identities

$$\Gamma(m - \delta) = \Gamma(-\delta) [-\delta]_m, \quad (2)$$

$$\Gamma(k + m - \delta) = \Gamma(m - \delta) [m - \delta]_k, \quad (3)$$

$$\Gamma(k + m - \delta + \gamma) = \Gamma(m - \delta + \gamma) [m - \delta + \gamma]_k, \quad (4)$$

then applying (2) ~ (4) into the RHS of (1), we obtain

$$\text{RHS of (1)} = e^{-i\pi\gamma} (z-b)^{\delta-\gamma}$$

$$\times \sum_{m=0}^{\infty} \frac{\Gamma(m-\delta+\gamma)}{m! \Gamma(-\delta)} \left(\frac{d}{z-b}\right)^m \sum_{k=0}^{\infty} \frac{[m-\delta+\gamma]_k}{k!} \left(\frac{c}{z-b}\right)^k \quad (5)$$

$$= e^{-i\pi\gamma} (z-b)^{\delta-\gamma} \sum_{m=0}^{\infty} \frac{\Gamma(m-\delta+\gamma)}{m! \Gamma(-\delta)} \left(\frac{d}{z-b}\right)^m \left(1 - \frac{c}{z-b}\right)^{\delta-\gamma-m} \quad (6)$$

$$= e^{-i\pi\gamma} (z-b-c)^{\delta-\gamma} \frac{\Gamma(-\delta+\gamma)}{\Gamma(-\delta)} \sum_{m=0}^{\infty} \frac{[-\delta+\gamma]_m}{m!} \left(\frac{d}{z-b-c}\right)^m \quad (7)$$

$$= e^{-i\pi\gamma} \frac{\Gamma(-\delta+\gamma)}{\Gamma(-\delta)} (z-b-c-d)^{\delta-\gamma}, \quad \left( \left| \frac{\Gamma(-\delta+\gamma)}{\Gamma(-\delta)} \right| < \infty \right) \quad (8)$$

using the relationship

$$\sum_{k=0}^{\infty} \frac{[\lambda]_k}{k!} z^k = (1-z)^{-\lambda}. \quad (9)$$

The result (8) is same as the one obtained by Lemma (i).

[II] When  $d=0$  and  $\delta=1$ , we obtain

$$(((z-b)^{\beta}-c)^{\alpha})_{\gamma} = e^{-i\pi\gamma} (z-b)^{\alpha\beta-\gamma}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-1]_m [-\alpha(1-m)]_k \Gamma(\beta k - \alpha\beta(1-m) + \gamma)}{m! \cdot k! \Gamma(\beta k - \alpha\beta(1-m))} \left(\frac{c}{(z-b)^{\beta}}\right)^k \left(\frac{0}{(z-b)^{\alpha\beta}}\right)^m \quad (10)$$

$$= e^{-i\pi\gamma} (z-b)^{\alpha\beta-\gamma} \sum_{k=0}^{\infty} \frac{[-\alpha]_k \Gamma(\beta k - \alpha\beta + \gamma)}{k! \Gamma(\beta k - \alpha\beta)} \left(\frac{c}{(z-b)^{\beta}}\right)^k \quad (11)$$

$$\left( \left| \frac{c}{(z-b)^{\beta}} \right| < 1, \quad \left| \frac{\Gamma(\beta k - \alpha\beta + \gamma)}{\Gamma(\beta k - \alpha\beta)} \right| < \infty \right)$$

from Theorem 1. (i).

That is, in this case we have Theorem D. (i) it self, clearly.

[III] When  $n=0$ , we have

$$(((z-b)^{\beta}-c)^{\alpha}-d)^{\delta})_0 = (z-b)^{\alpha\beta\delta}$$

$$\times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k}{m! \cdot k!} \left(\frac{c}{(z-b)^{\beta}}\right)^k \left(\frac{d}{(z-b)^{\alpha\beta}}\right)^m. \quad (12)$$

$$((z-b)^\beta - c)^\alpha - d \neq 0, \quad \left| \frac{c}{(z-b)^\beta} \right| < 1, \quad \left| \frac{d}{(z-b)^{\alpha\beta}} \right| < 1, .$$

from Theorem 1. (ii).

Indeed we obtain

$$\text{RHS of (12)} = (z-b)^{\alpha\beta\delta}$$

$$\times \sum_{m=0}^{\infty} \frac{[-\delta]_m}{m!} \left( \frac{d}{(z-b)^{\alpha\beta}} \right)^m \sum_{k=0}^{\infty} \frac{[-\alpha(\delta-m)]_k}{k!} \left( \frac{c}{(z-b)^\beta} \right)^k \quad (13)$$

$$= (z-b)^{\alpha\beta\delta} \sum_{m=0}^{\infty} \frac{[-\delta]_m}{m!} \left( \frac{d}{(z-b)^{\alpha\beta}} \right)^m \left( 1 - \frac{c}{(z-b)^\beta} \right)^{\alpha\delta - \alpha m} \quad (14)$$

$$= ((z-b)^\beta - c)^{\alpha\delta} \sum_{m=0}^{\infty} \frac{[-\delta]_m}{m!} \left( \frac{d}{((z-b)^\beta - c)^\alpha} \right)^m \quad (15)$$

$$= ((z-b)^\beta - c)^{\alpha\delta} \left( 1 - \frac{d}{((z-b)^\beta - c)^\alpha} \right)^\delta \quad (16)$$

$$= (((z-b)^\beta - c)^\alpha - d)^\delta, \quad (17)$$

clearly.

[IV] When  $n=1$ , we have

$$\begin{aligned} & (((z-b)^\beta - c)^\alpha - d)^\delta \Big|_1 = -(z-b)^{\alpha\beta\delta-1} \\ & \times \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k \cdot (\beta k - \alpha\beta\delta + \alpha\beta m)}{m! \cdot k!} \left( \frac{c}{(z-b)^\beta} \right)^k \left( \frac{d}{(z-b)^{\alpha\beta}} \right)^m, \quad (18) \end{aligned}$$

from Theorem 1. (ii).

Then letting

$$R := \left( \frac{u}{(z-b)^\beta} \right)^\alpha \quad (u = (z-b)^\beta - c) \quad (19)$$

and

$$S := \left( 1 - \frac{d}{u^\alpha} \right)^\delta \quad (20)$$



we have

$$\sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k \cdot \beta k}{m! \cdot k!} \left( \frac{c}{(z-b)^\beta} \right)^k \left( \frac{d}{(z-b)^{\alpha\beta}} \right)^m$$

$$= \frac{-c\alpha\beta\delta}{u} R^\delta S - \frac{cd\alpha\beta\delta}{u(u^\alpha-d)} R^\delta S, \quad (21)$$

$$- \alpha\beta\delta \sum_{m,k=0}^{\infty} \frac{[-\delta]_m [-\alpha(\delta-m)]_k}{m! \cdot k!} \left( \frac{c}{(z-b)^\beta} \right)^k \left( \frac{d}{(z-b)^{\alpha\beta}} \right)^m$$

$$= -\alpha\beta\delta R^\delta S, \quad (22)$$

and

$$\alpha\beta \sum_{m,k=0}^{\infty} \frac{[-\delta]_m \cdot m [-\alpha(\delta-m)]_k}{m! \cdot k!} \left( \frac{c}{(z-b)^\beta} \right)^k \left( \frac{d}{(z-b)^{\alpha\beta}} \right)^m$$

$$= -\frac{d\alpha\beta\delta}{(u^\alpha-d)} R^\delta S, \quad (23)$$

Therefore, applying (23), (22) and (21) into (18), we obtain

$$(((z-b)^\beta - c)^\alpha - d)^\delta)_1 = (z-b)^{\alpha\beta\delta-1} R^\delta S$$

$$\times \left[ \frac{c\alpha\beta\delta}{u} + \frac{cd\alpha\beta\delta}{u(u^\alpha-d)} + \alpha\beta\delta + \frac{d\alpha\beta\delta}{u^\alpha-d} \right] \quad (24)$$

$$= \alpha\beta\delta (z-b)^{\alpha\beta\delta-1} \left( \frac{c}{u} + 1 \right) \left( \frac{d}{u^\alpha-d} + 1 \right) \quad (25)$$

$$= \alpha\beta\delta (z-b)^{-1} (u^\alpha-d)^\delta \left( \frac{(z-b)^\beta}{u} \right) \left( \frac{u^\alpha}{u^\alpha-d} \right) \quad (26)$$

$$= \alpha\beta\delta (u^\alpha-d)^{\delta-1} (u^{\alpha-1}) (z-b)^{\beta-1} \quad (27)$$

$$= \alpha\beta\delta (((z-b)^\beta - c)^\alpha - d)^{\delta-1} ((z-b)^\beta - c)^{\alpha-1} (z-b)^{\beta-1} \quad (28)$$

clearly.

## References

- [1] K. Nishimoto ; Fractional Calculus, Vol. 1 (1984), Vol. 2 (1987), Vol. 3 (1989), Vol. 4 (1991), Vol. 5, (1996), Descartes Press, Koriyama, Japan.
- [2] K. Nishimoto ; An Essence of Nishimoto's Fractional Calculus (Calculus of the 21st Century); Integrals and Differentiations of Arbitrary Order (1991), Descartes Press, Koriyama, Japan.
- [3] K. Nishimoto ; On Nishimoto's fractional calculus operator  $N^\nu$  ( On an action group ), J. Frac. Calc. Vol. 4, Nov. (1993), 1 - 11.
- [4] K. Nishimoto ; Unification of the integrals and derivatives ( A serendipity in fractional calculus ), J. Frac. Calc. Vol. 6, Nov. (1994), 1 - 14.
- [5] K. Nishimoto ; Ring and Field Produced from The Set of N- Fractional Calculus Operator, J. Frac Calc. Vol. 24, Nov. (2003), 29 - 36.
- [6] K. Nishimoto; On the fractional calculus of functions  $(a - z)^\beta$  and  $\log(a - z)$ , J. Frac. Calc. Vol.3, May (1993), 19 - 27.
- [7] K. Nishimoto and Shih- Tong Tu ; Fractional calculus of Psi functions ( Generalized Polygamma functions ), J. Frac. Calc. Vol.5, May (1994), 27 -34.
- [8] Shih- Tong Tu and K. Nishimoto ; On the fractional calculus of functions  $(cz - a)^\beta$  and  $\log(cz - a)$ , J. Frac.Calc.Vol.5, May (1994), 35 - 43.
- [9] K. Nishimoto ; N- Fractional Calculus of the Power and Logarithmic Functions, and Some Identities , J. Frac. Calc. Vol.21, May (2002), 1 - 6.
- [10] K. Nishimoto ; Some Theorems for N- Fractional Calculus of Logarithmic Functions I, J. Frac Calc.Vol. 21, May (2002), 7 - 12.
- [11] K. Nishimoto ; N- Fractional Calculus of Products of Some Power Functions, J. Frac. Calc.Vol.27, May (2005), 83 - 88..
- [12] K. Nishimoto ; N- Fractional Calculus of Some Composite Functions, J. Frac. Calc. Vol. 29, May (2006), 35 - 44.
- [13] K. Nishimoto ; N- Fractional Calculus of Some Elementary Functions and Their Semi Differintegrations, J. Frac. Calc. Vol. 30, Nov. (2006), 1- 10.
- [14] David Dummit and Richard M. Foote ; Abstract Algebra, Prentice Hall (1991).
- [15] K.B. Oldham and J. Spanier ; The Fractional Calculus, Academic Press (1974).
- [16] A.C. McBride ; Fractional Calculus and Integral Transformations of Generalized Functions, Research Notes, Vol. 31, (1979), Pitman.
- [17] S. G. Samko, A.A. Kilbas and O.I. Marichev ; Fractional Integrals and Derivatives, and Some Their Applications (1987), Nauka, USSR.
- [18] K.S. Miller and B. Ross ; An Introduction to The Fractional Calculus and Fractional Diffetial Equations, John Wiley & Sons, (1993).
- [19] V. Kiryakova ; Generalized fractional calculus and applications, Pitman Research Notes, No. 301, (1994), Longman.
- [20] Igor Podlubny ; Fractional Differential Equations (1999), Academic Press.
- [21] R. Hilfer (Ed.) ; Applications of Fractional Calculus in Physics, (2000), World Scientific, Singapor, New Jersey, London, Hong Kong.
- [22] A.P. Prudnikov, Yu. A. Bryckov and O.I. Marichev ; Integrals and Series, Vol. I, Gordon and Breach, New York, (1986).
- [23] S. Moriguchi, K. Udagawa and S. Hitotsumatsu ; Mathematical Formulae, Vol. 2, Iwanami Zensho, (1957), Iwanami, Japan.

Katsuyuki Nishimoto  
 Institute for Applied Mathematics  
 Descartes Press Co.  
 2 - 13 - 10 Kaguike, Koriyama  
 963 - 8833 Japan